



Rio Dell City Hall
675 Wildwood Avenue
Rio Dell, CA 95562
(707) 764-3532
cityofriodell.ca.gov

PRESS RELEASE

12-21-2020

Rio Dell Observes Spike in COVID-19

Since October 1st of this year, the City of Rio Dell has been conducting COVID-19 surveillance testing at its Wastewater Treatment Plant. The testing, through Biobot Analytics, samples the inflow of untreated wastewater in order to isolate the unique genetic signature of SARS-CoV-2 (COVID-19) and analyze the amount of the virus present. The Biobot Analytics testing is done independently and results have been forwarded to the Public Health Officer since the inception of the program. Between October 1st and December 3rd test results have ranged from 'non-detect' to detection at levels below the national average.

On December 17, 2020 the City of Rio Dell conducted its latest test showing higher concentration levels than 84% of all quantifiable samples collected nationwide by Biobot Analytics in the past six weeks. This level is well above the median national average.

"This information further supplements and confirms the observations of the County Health Department." Stated Rio Dell City Manager Kyle Knopp. "The virus is present in this small community at high levels. We provide this information as another tool residents can use when making their plans over the holiday season and near future."

The City continues to encourage mask wearing, social distancing, regular hand washing, regular surface cleaning, testing and staying home when sick as good methods to slow the virus and protect the community's fragile health systems.

Rio Dell residents have nearby access to testing:

Palco Pharmacy (appointment required)
113 Main St Suite D, Scotia, CA 95565
(707) 764-3591

Southern Trinity Health Services, Scotia Bluffs (call first)
500 B Street, Scotia, California 95565
(707) 764-5617

Optum Serve
Redwood Acres and occasionally Fortuna (register beforehand)
lhi.care
888-634-1123

///



BIO
BOT

BIOBOT ANALYTICS

Pioneering a better way to track Covid-19

The nation's leader in wastewater epidemiology

SARS-CoV-2 is shed in the stool of Covid-19 patients, making its way into our sewers. We analyze sewage to determine the presence of infected individuals and estimate the number of cases.

Trend analysis

Determine when to safely re-open, and keep open, our communities and businesses through frequent sampling.

Early warning

Detect the re-emergence of Covid-19 in a population to rapidly take action and contain new outbreaks.

Test at scale

For a fraction of the cost of mass individual testing, get an overview of the scope of the outbreak.

Comprehensive product

1. **Program design.** We develop the appropriate sampling program for your needs.
2. **Sample collection.** We provide sampling protocols, mail-in kits, and hardware.
3. **Lab analysis.** We isolate the unique genetic signature of SARS-CoV-2 and analyze the amount of the virus present.
4. **Data analysis.** Leveraging our data science models, built with thousands of samples, we estimate the prevalence of Covid-19 in the sample population.
5. **Data interpretation.** Through data visualizations and weekly webinars, we provide expert customer support.
6. **Additional product lines.** Be first to access our additional product lines: opioids, antibiotic resistant bacteria, and others under development.


Our Covid-19 testing presence

400 testing sites

42 states represented

~10% of the U.S. population

Weekly testing



Contact Us

covid19@biobot.io
www.biobot.io
@BiobotAnalytics

City of Rio Dell WWTF

Sample collection date: **December 17, 2020**

SARS-CoV-2 virus in sewage

DETECTED

Virus concentration
(genome copies per liter of sewage)


616,146

Normalized* virus concentration
(genome copies per liter of sewage)

983,901

**Normalized virus concentration value is derived by adjusting the raw virus concentration to the PMMV fecal marker in order to account for dilution*

Normalized virus concentration over time



Regional overview of normalized virus concentration levels - West


State-level mean of samples
(collected in the past 6 weeks)

lower higher


Color indicates normalized virus concentration level

Your state outlined in red

Size of circle indicates population served by participating facilities, ranging from 10k to 1mil




Normalized virus concentrations trend with new cases



Normalized virus concentrations in comparison

Your sample has higher concentration levels than **84%** of all quantifiable samples collected in the past six weeks.



Biobot COVID19 incidence estimate*

* using Biobot's current analysis model v2.0, which reflects active R&D and will change over time with developing research

1 to 5 new cases / day
(0.11% incidence rate)

Using a reported flow rate of 0.37 MGD

For more information, read the whitepaper:
<https://doi.org/10.1101/2020.06.15.20117747>

This incidence estimate represents the projected average of **confirmed new clinical cases (per day)** that will be reported in your community 7 days from the sampling date. This estimate reflects active R&D.

This number is derived from Biobot's latest proprietary case model, leveraging thousands of samples analyzed for Covid-19. For context, USAFacts reports **17 new cases** on this sampling date and an average of **25.6 cases** per day for the past 7 days in **Humboldt County, CA**.

Visit our website for more details behind the process:
<https://www.biobot.io/case-estimates>
<https://www.biobot.io/updated-model>



Biobot's Covid-19 wastewater testing product

Lab protocol

Our methods for detecting SARS-CoV-2 in sewage are adapted from CDC protocols. Our approach relies on detecting genetic fragments of the virus that are excreted in stool by qPCR analysis, which does not determine if the virus is dead or active.

Limit of detection (LOD)

The LOD for our lab protocol is 3,600 copies/L of sewage (see more details in Release Notes below). In terms of case estimates, we reliably detect the virus (>99%) when there is at least 1 infected person in a population of 6,500 people.

Data use

Biobot's wastewater data provides an alternative metric to guide response to the Covid-19 outbreak. We recommend sharing this information with local public health officials. We believe this work has the greatest impact on a statewide level, and hope that you will reach out to your local officials and encourage the expansion of our partnership across your state.

Questions and support

For questions specific to your report, email support@biobot.io.

Biobot's QA/QC protocol

Biobot has an in-house lab facility with a team of scientists dedicated 100% to Covid-19 wastewater testing. All reported data passes our QA/QC protocol:

1) Sample collection

- 3 x 50 ml samples are shipped with a frozen pack to keep 4C temperature control.
- Documentation collected via online form: location, date, time, flow rate on sampling day, sampling type, precipitation events.

2) Storage

- Raw sewage samples are received at Biobot and immediately pasteurized. Pasteurized samples are stored at 4C for up to 3 days before viral concentration.
- Extracted RNA is stored at 4C for no longer than 24h before analysis by RT-qPCR.
- Extracted RNA is stored at -80C for the next 12 months.

3) Sample processing

- 15 mL of sewage sample is used for viral concentration and RNA extraction.
- Second and third replicates are kept at 4C for 30 days as back-up.
- Pepper Mild Mottle Virus, PMMoV, is a fecal indicator used as internal control.
- CDC Primers N1 and N2 are used to target SARS-CoV-2.
- Each test primer (N1 and N2) is run in triplicate in the qPCR assay.
- Four positive controls (synthetic SARS-CoV-2 N gene) are run in each 96-well plate.
- Two negative controls (no template) are run in each 96-well plate.
- Standard curves (synthetic SARS-CoV-2 N gene) are run once a week.

Biobot's data interpretation

Raw viral concentration (genome copies per L of sewage)

The raw SARS-CoV-2 viral concentration is directly measured by the laboratory qPCR assay.



Normalized viral concentration (genome copies per L of sewage)

We normalize the SARS-CoV-2 viral concentration to a fecal indicator, to account for differences in dilution. We use PMMV as this fecal indicator, which is an RNA virus that is commonly excreted in stool.

Biobot's Covid-19 incidence estimate

Our latest Covid-19 incidence estimation model is built from Biobot's dataset, the largest Covid-19 dataset in the world to date. We mined this dataset to derive an empirical relationship between the amount of virus in sewage samples and the number cases reported in the associated communities over the next 7 days. This means that our model provides an estimate of the number of cases that will be reported in your community in the next week.

We convert the raw viral concentration into a Covid-19 case estimate using the following equation:

$$\text{Number of Covid19 cases} = \frac{\text{Viral concentration} * \text{Flow rate on sampling date}}{\text{Virus shed per reported case per day}^{**}}$$

* We use the raw viral concentration in this calculation because dilution is accounted for by multiplying the raw concentration with the daily flow.

** Parameter currently derived from Biobot's dataset. Learn more at: <https://www.biobot.io/updated-model>

The incidence estimation equation uses your reported flow rate and the measured virus concentration, but does not use your reported catchment population. If we do not have a reported flow rate for your sample, we use your location's average influent flow rate that you provided during onboarding in the calculation.

You can interpret this incidence estimate as the number of cases that will be reported in your community in the next 4-7 days. For more information on the predictive nature of wastewater data, read our blog post: <https://www.biobot.io/case-estimates>. For more information on our new incidence estimation model, read our blog post: <https://www.biobot.io/updated-model>. For more technical details on both aspects, read the preprint: <https://doi.org/10.1101/2020.06.15.20117747>.

Our model is built in part on data from clinically confirmed cases. Reported cases include only patients who sought out Covid-19 testing and received a positive test result. This population likely does not include most asymptomatic patients or individuals without access to testing. Because reported cases are an undercount of the true number of infected individuals, our estimate is also likely a conservative projection of the number of new cases in your population. We are working actively to update our model based on the latest science and data sets available.

Incidence rate (%)

The incidence rate is calculated by dividing the Biobot case estimate by your reported catchment population.

Release notes

Lab protocol versions

We are continuously working to improve our protocols to increase the sensitivity of our measurements and reduce variability. You can find the protocol that was used to generate your data at the bottom of each page of this report:

Lab Protocol Version	Limit of detection (LOD)	Description
v2.3 (current)	3,600 copies/L	Kit-based virus concentration and RNA extraction with one-step RT-qPCR at Biobot and an improved algorithmic Ct calling method.
v2.2	2,100 copies/L	Kit-based virus concentration and RNA extraction with one-step RT-qPCR at Biobot laboratory.
v2.1	1,700 copies/L	Kit-based virus concentration and RNA extraction with one-step RT-qPCR at our MIT partner laboratory.
v2.0	34,000 copies/L	Kit-based virus concentration and RNA extraction with two-step RT-qPCR at our MIT partner laboratory.
v1.0	6,400 copies/L	PEG virus concentration and Trizol RNA extraction with two-step RT-qPCR at our MIT partner laboratory.

Data analysis & model versions

We are constantly iterating on and improving our data processing, analysis, and Covid-19 models to improve the interpretability of our data. You can see which version of our analysis and model was used in this report at the bottom of each page, and you can find more specific details in the release notes below.

Data Analysis & Model version	Description
v2.0 (current)	We launched an updated case estimation model which uses a viral shedding parameter derived from mining our proprietary wastewater dataset. The basic equation is the same as before, but the “viral shedding” parameter is now empirically derived as the amount of virus shed per reported case (rather than based on clinical studies of shedding in stool). As for previous versions, the case estimate is calculated using the raw SARS-CoV-2 concentration, and all other plots are generated using the normalized concentration, normalized with the same method as prior versions.
v1.2	We updated our normalization process for the virus concentration to retain units of copies/L of sewage. We multiply the raw lab concentration by a scaling factor ($scaling\ factor = \frac{reference\ PMMV}{kit\ PMMV}$). The reference PMMV is derived empirically from our entire database. As in previous versions, the case estimate is calculated using the raw SARS-CoV-2 concentration and accounts for dilution by using the flow rate provided.
v1.1	We updated detection thresholds to reduce the chance of false positives. Specifically, we’ve raised our limit of detection to ensure that all measurements can be confidently quantified, and are requiring two positive measurements per sample (out of six) to consider a sample detected.

Data Analysis & Model version	Description
v1.0	Raw viral concentration and Covid-19 case estimates are reported. The model parameter (virus shed per infected person per day) is determined from direct communications with Professor Kyle Bibby and Dr. Aaron Bivins and based on clinical viral shedding reported in Wolfel et al. <i>Nature</i> (2020).

Report design versions

We are continually making updates to our report based on internal R&D and feedback from our customers.


Report Design version	Description
v3.4 (current)	Major changes: - Implemented 7-day rolling average and daily cases in county for time series comparison plot. - Included comparison to historical 7-day average of reported cases in case estimate box and scatter plot.
v3.3.1	Minor changes: Show Maryland and Delaware in Northeast map instead of South.
v3.3	Major changes: - All plots with aggregate data analyze samples based on a six week rolling window up to and including the date the sample was collected. Minor changes: - Added visual indicators to timeseries plot for new cases reported on the sampling date. - Regional geographic maps display the mean of the samples taken during the six week window, instead of the median.
v3.2.3	Minor changes: Added the external Kit ID reference in addition to Biobot's internal Kit ID.
v3.2.2	Minor changes: - Updated the case estimate title to incidence estimate to better reflect the interpretation of our current model. - Changed the incidence estimate description to better reflect the updated model, and provided details around the methodology of how we arrived at the new model.



Report Design version	Description
v3.2.1	Minor changes: • The case estimate is provided with an updated viral shedding parameter, and reflects the number of cases which will be reported in your community next week. • We've included information about your county's reported new cases back into the report.
v3.2	Major changes: • The normalized viral concentration is shown in the report, so you can reproduce the time series plot on your own. • The sampling date shows the sampling end date (before it was the start date) to be in line with the industry standard. • We removed the box with county-level data from USA Facts on cumulative and new cases, so you can compare the Biobot data with your most relevant local datasets. Minor changes: • We changed the language around case estimates based on internal R&D to reflect new cases that will be predicted in your county by next week. • We updated language on charts to reflect the new normalization units, and removed redundant explanatory text. • We added legends to the timeseries and sample distribution plots. • We updated all visualizations to reflect data collected since June. • We removed the "percent change" callout in the time series and added axes with units.

City of Rio Dell WWTFSample collection date: **December 3, 2020****SARS-CoV-2 virus in sewage****DETECTED**Virus concentration
(genome copies per liter of sewage)**6,215**Normalized* virus concentration
(genome copies per liter of sewage)**4,705**

**Normalized virus concentration value is derived by adjusting the raw virus concentration to the PMMV fecal marker in order to account for dilution*


Normalized virus concentration over time**Regional overview of normalized virus concentration levels - West**State-level mean of samples
(collected in the past 6 weeks)

lower higher


Color indicates normalized virus concentration level

Your state outlined in red

Size of circle indicates population served by participating facilities, ranging from 10k to 1mil




Normalized virus concentrations trend with new cases



Normalized virus concentrations in comparison

Your sample has higher concentration levels than **1%** of all quantifiable samples collected in the past six weeks.



Biobot COVID19 incidence estimate*

* using Biobot's current analysis model v2.0, which reflects active R&D and will change over time with developing research

< 1 new cases / day
(0.0007% incidence rate)

Using a reported flow rate of 0.22 MGD

For more information, read the whitepaper:
<https://doi.org/10.1101/2020.06.15.20117747>

This incidence estimate represents the projected average of **confirmed new clinical cases (per day)** that will be reported in your community 7 days from the sampling date. This estimate reflects active R&D.

This number is derived from Biobot's latest proprietary case model, leveraging thousands of samples analyzed for Covid-19. For context, USAFacts reports **44 new cases** on this sampling date and an average of **24.9 cases** per day for the past 7 days in **Humboldt County, CA**.

Visit our website for more details behind the process:
<https://www.biobot.io/case-estimates>
<https://www.biobot.io/updated-model>



Biobot's Covid-19 wastewater testing product

Lab protocol

Our methods for detecting SARS-CoV-2 in sewage are adapted from CDC protocols. Our approach relies on detecting genetic fragments of the virus that are excreted in stool by qPCR analysis, which does not determine if the virus is dead or active.

Limit of detection (LOD)

The LOD for our lab protocol is 3,600 copies/L of sewage (see more details in Release Notes below). In terms of case estimates, we reliably detect the virus (>99%) when there is at least 1 infected person in a population of 6,500 people.

Data use

Biobot's wastewater data provides an alternative metric to guide response to the Covid-19 outbreak. We recommend sharing this information with local public health officials. We believe this work has the greatest impact on a statewide level, and hope that you will reach out to your local officials and encourage the expansion of our partnership across your state.

Questions and support

For questions specific to your report, email support@biobot.io.

Biobot's QA/QC protocol

Biobot has an in-house lab facility with a team of scientists dedicated 100% to Covid-19 wastewater testing. All reported data passes our QA/QC protocol:

1) Sample collection

- 3 x 50 ml samples are shipped with a frozen pack to keep 4C temperature control.
- Documentation collected via online form: location, date, time, flow rate on sampling day, sampling type, precipitation events.

2) Storage

- Raw sewage samples are received at Biobot and immediately pasteurized. Pasteurized samples are stored at 4C for up to 3 days before viral concentration.
- Extracted RNA is stored at 4C for no longer than 24h before analysis by RT-qPCR.
- Extracted RNA is stored at -80C for the next 12 months.

3) Sample processing

- 15 mL of sewage sample is used for viral concentration and RNA extraction.
- Second and third replicates are kept at 4C for 30 days as back-up.
- Pepper Mild Mottle Virus, PMMoV, is a fecal indicator used as internal control.
- CDC Primers N1 and N2 are used to target SARS-CoV-2.
- Each test primer (N1 and N2) is run in triplicate in the qPCR assay.
- Four positive controls (synthetic SARS-CoV-2 N gene) are run in each 96-well plate.
- Two negative controls (no template) are run in each 96-well plate.
- Standard curves (synthetic SARS-CoV-2 N gene) are run once a week.

Biobot's data interpretation

Raw viral concentration (genome copies per L of sewage)

The raw SARS-CoV-2 viral concentration is directly measured by the laboratory qPCR assay.



Normalized viral concentration (genome copies per L of sewage)

We normalize the SARS-CoV-2 viral concentration to a fecal indicator, to account for differences in dilution. We use PMMV as this fecal indicator, which is an RNA virus that is commonly excreted in stool.

Biobot's Covid-19 incidence estimate

Our latest Covid-19 incidence estimation model is built from Biobot's dataset, the largest Covid-19 dataset in the world to date. We mined this dataset to derive an empirical relationship between the amount of virus in sewage samples and the number cases reported in the associated communities over the next 7 days. This means that our model provides an estimate of the number of cases that will be reported in your community in the next week.

We convert the raw viral concentration into a Covid-19 case estimate using the following equation:

$$\text{Number of Covid19 cases} = \frac{\text{Viral concentration} * \text{Flow rate on sampling date}}{\text{Virus shed per reported case per day}^{**}}$$

* We use the raw viral concentration in this calculation because dilution is accounted for by multiplying the raw concentration with the daily flow.

** Parameter currently derived from Biobot's dataset. Learn more at: <https://www.biobot.io/updated-model>

The incidence estimation equation uses your reported flow rate and the measured virus concentration, but does not use your reported catchment population. If we do not have a reported flow rate for your sample, we use your location's average influent flow rate that you provided during onboarding in the calculation.

You can interpret this incidence estimate as the number of cases that will be reported in your community in the next 4-7 days. For more information on the predictive nature of wastewater data, read our blog post: <https://www.biobot.io/case-estimates>. For more information on our new incidence estimation model, read our blog post: <https://www.biobot.io/updated-model>. For more technical details on both aspects, read the preprint: <https://doi.org/10.1101/2020.06.15.20117747>.

Our model is built in part on data from clinically confirmed cases. Reported cases include only patients who sought out Covid-19 testing and received a positive test result. This population likely does not include most asymptomatic patients or individuals without access to testing. Because reported cases are an undercount of the true number of infected individuals, our estimate is also likely a conservative projection of the number of new cases in your population. We are working actively to update our model based on the latest science and data sets available.

Incidence rate (%)

The incidence rate is calculated by dividing the Biobot case estimate by your reported catchment population.

Release notes

Lab protocol versions

We are continuously working to improve our protocols to increase the sensitivity of our measurements and reduce variability. You can find the protocol that was used to generate your data at the bottom of each page of this report:

Lab Protocol Version	Limit of detection (LOD)	Description
v2.3 (current)	3,600 copies/L	Kit-based virus concentration and RNA extraction with one-step RT-qPCR at Biobot and an improved algorithmic Ct calling method.
v2.2	2,100 copies/L	Kit-based virus concentration and RNA extraction with one-step RT-qPCR at Biobot laboratory.
v2.1	1,700 copies/L	Kit-based virus concentration and RNA extraction with one-step RT-qPCR at our MIT partner laboratory.
v2.0	34,000 copies/L	Kit-based virus concentration and RNA extraction with two-step RT-qPCR at our MIT partner laboratory.
v1.0	6,400 copies/L	PEG virus concentration and Trizol RNA extraction with two-step RT-qPCR at our MIT partner laboratory.

Data analysis & model versions

We are constantly iterating on and improving our data processing, analysis, and Covid-19 models to improve the interpretability of our data. You can see which version of our analysis and model was used in this report at the bottom of each page, and you can find more specific details in the release notes below.

Data Analysis & Model version	Description
v2.0 (current)	We launched an updated case estimation model which uses a viral shedding parameter derived from mining our proprietary wastewater dataset. The basic equation is the same as before, but the “viral shedding” parameter is now empirically derived as the amount of virus shed per reported case (rather than based on clinical studies of shedding in stool). As for previous versions, the case estimate is calculated using the raw SARS-CoV-2 concentration, and all other plots are generated using the normalized concentration, normalized with the same method as prior versions.
v1.2	We updated our normalization process for the virus concentration to retain units of copies/L of sewage. We multiply the raw lab concentration by a scaling factor ($scaling\ factor = \frac{reference\ PMMV}{kit\ PMMV}$). The reference PMMV is derived empirically from our entire database. As in previous versions, the case estimate is calculated using the raw SARS-CoV-2 concentration and accounts for dilution by using the flow rate provided.
v1.1	We updated detection thresholds to reduce the chance of false positives. Specifically, we’ve raised our limit of detection to ensure that all measurements can be confidently quantified, and are requiring two positive measurements per sample (out of six) to consider a sample detected.

Data Analysis & Model version	Description
v1.0	Raw viral concentration and Covid-19 case estimates are reported. The model parameter (virus shed per infected person per day) is determined from direct communications with Professor Kyle Bibby and Dr. Aaron Bivins and based on clinical viral shedding reported in Wolfel et al. <i>Nature</i> (2020).

Report design versions

We are continually making updates to our report based on internal R&D and feedback from our customers.

Report Design version	Description
v3.4 (current)	Major changes: - Implemented 7-day rolling average and daily cases in county for time series comparison plot. - Included comparison to historical 7-day average of reported cases in case estimate box and scatter plot.
v3.3.1	Minor changes: Show Maryland and Delaware in Northeast map instead of South.
v3.3	Major changes: - All plots with aggregate data analyze samples based on a six week rolling window up to and including the date the sample was collected. Minor changes: - Added visual indicators to timeseries plot for new cases reported on the sampling date. - Regional geographic maps display the mean of the samples taken during the six week window, instead of the median.
v3.2.3	Minor changes: Added the external Kit ID reference in addition to Biobot's internal Kit ID.
v3.2.2	Minor changes: - Updated the case estimate title to incidence estimate to better reflect the interpretation of our current model. - Changed the incidence estimate description to better reflect the updated model, and provided details around the methodology of how we arrived at the new model.



Report Design version	Description
v3.2.1	Minor changes: • The case estimate is provided with an updated viral shedding parameter, and reflects the number of cases which will be reported in your community next week. • We've included information about your county's reported new cases back into the report.
v3.2	Major changes: • The normalized viral concentration is shown in the report, so you can reproduce the time series plot on your own. • The sampling date shows the sampling end date (before it was the start date) to be in line with the industry standard. • We removed the box with county-level data from USA Facts on cumulative and new cases, so you can compare the Biobot data with your most relevant local datasets. Minor changes: • We changed the language around case estimates based on internal R&D to reflect new cases that will be predicted in your county by next week. • We updated language on charts to reflect the new normalization units, and removed redundant explanatory text. • We added legends to the timeseries and sample distribution plots. • We updated all visualizations to reflect data collected since June. • We removed the "percent change" callout in the time series and added axes with units.